

High-energy gamma-rays from GRB X-ray flares

X. Y. Wang^{*,†}, Z. Li^{**} and P. Mészáros^{*}

^{*}*Department of Astronomy and Astrophysics, Penn. State University, University Park, PA 16802*

[†]*Department of Astronomy, Nanjing University, Nanjing 210093, China*

^{**}*Physics Faculty, Weizmann Institute of Science, Rehovot 76100, Israel*

Abstract. The recent detection of X-ray flares during the afterglow phase of gamma-ray bursts (GRBs) suggests an inner-engine origin, at radii inside the forward shock. There must be inverse Compton (IC) emission arising from such flare photons scattered by forward shock afterglow electrons when they are passing through the forward shock. We find that this IC emission produces high energy gamma-ray flares, which may be detected by AGILE, GLAST and ground-based TeV telescopes. The anisotropic IC scattering between flare photons and forward shock electrons does not affect the total IC component intensity, but cause a time delay of the IC component peak relative to the flare peak. The anisotropic scattering effect may also weaken, to some extent, the suppression effect of the afterglow intensity induced by the enhanced electron cooling due to flare photons. We speculate that this IC component may already have been detected by EGRET from a very strong burst—GRB940217. Future observations by GLAST may help to distinguish whether X-ray flares originate from late central engine activity or from external shocks.

IC OF FLARE PHOTONS BY FORWARD SHOCK ELECTRONS

X-ray flares have been detected during the early afterglows in a significant fraction of gamma-ray bursts (e.g. [1]; [4]; [5]). The amplitude of the X-ray flare can be a factor of up to ~ 500 for GRB050502B [1, 3], and in most cases a factor several, compared with the background afterglow component. The rapid rise and decay behavior of some flares suggests that they are caused by internal dissipation of energy due to late central engine activity. Therefore the inner flare photons, when they are passing through the forward shocks, must interact with the shocked electrons and get boosted to higher energies [6]. See Fig.1 for a cartoon illustration. The observed overlapping in time between flares and afterglows indicates that these two kind of particles (i.e. photons and shocked electrons) have indeed interacted with each other around the overlap time[2]. This process is expected to occur whether the X-ray flares are produced by internal shocks or by other internal energy dissipation mechanisms, as long as they occur inside the inner edge of the forward shock region.

Let us now estimate the flux of this IC component. We consider an X-ray flare of duration δt superimposed upon an underlying power law X-ray afterglow around time $t = 10^3 t_3$ s after the burst, as observed in GRB050502B. We find that the energy density of flare photons in the comoving frame of the forward shock is larger than the magnetic energy density ($B^2/8\pi$) when $F_X > 10^{-10} \varepsilon_{B,-2} E_{52} t_3^{-1} D_{28}^{-2} \text{ erg cm}^{-2} \text{ s}^{-1}$, where ε_B is the equipartition factor for magnetic field in forward shocks. Since the flux of the X-ray flare is also much larger than that of the underlying X-ray afterglow which may represent the synchrotron luminosity, we conclude that generally the




FIGURE 1. Cartoon illustration of the IC scattering between flare photons and forward shock electrons.

cooling of forward shock electrons is dominated by scattering the X-ray flare photons. It is also shown that when the X-ray flare flux is larger than a critical flux $F_{X,c} = 3 \times 10^{-10} E_{52}^{1/2} \epsilon_{e,-1}^{-1} n_0^{-1/2} t_3^{-1/2} D_{28}^{-2} \text{ergs s}^{-1} \text{cm}^{-2}$, we have $\gamma_m > \gamma_c$, and all the newly shocked electrons will cool, emitting most of their energy into the IC emission[6]. This critical flux is usually much lower than the X-ray flare flux averaged over its duration, so we conclude that the flare photons can effectively cool the electrons in the forward shock and most energy of the newly shock electrons will be lost into IC emission. So the total IC energy is $E_{IC} = \delta t L_e = 2 \times 10^{51} \epsilon_{e,-1} E_{52} (\delta t / t_p) \text{erg}$ and the averaged IC flux is

$$F_{IC} \simeq 10^{-9} \epsilon_{e,-1} E_{52} t_{p,3}^{-1} D_{28}^{-2} (\delta t / t_p) \text{ergs s}^{-1} \text{cm}^{-2}, \quad (1)$$

where $\delta t < t = t_p$ and t_p denotes the peak time of the flare [6]. The observed IC νF_ν flux peaks at

$$\epsilon_{IC,p} \simeq 2 \gamma_m^2 \epsilon_X \simeq 3 \epsilon_{e,-1}^2 E_{52}^{1/4} n_0^{-1/4} t_3^{-3/4} \epsilon_{X,\text{keV}} \text{GeV} \quad (2)$$

where ϵ_X is the peak of the flare energy spectrum. The IC emission occurs right inside the GLAST window and has a total fluence about $10^{-7} - 10^{-6} \epsilon_{e,-1} E_{52} D_{28}^{-2} \text{erg cm}^{-2}$ for $0.1 < \delta t / t < 1$, so it should be detected by GLAST. For the strongest bursts with $E \sim 10^{54} \text{erg}$, the GeV photons could even have been detected by EGRET, such as from GRB940217 [10]. AGILE may also be able to detect this high energy component from strong bursts.

If described approximately as broken power law, the IC energy spectrum (νF_ν) has indices of $1/2$ and $-(p-2)/2$ before and after the break at $\epsilon_{IC,p}$ respectively. The $-(p-2)/2$ power law spectrum can extend to a maximum energy $\epsilon_{IC,M}$, above which the IC falls into the Klein-Nishina regime. Requiring the IC scattering to be in the Thomson regime, $\gamma_e < \gamma_{e,M} = \Gamma m_e c^2 / \epsilon_X$, gives $\epsilon_{IC,M} = 0.4 E_{52}^{1/4} n_0^{-1/4} t_3^{-3/4} \epsilon_{X,\text{keV}}^{-1} \text{TeV}$. The optical depth due to $\gamma\gamma$ absorption on the X-ray flare photons for the maximum energy $\epsilon_{IC,M}$ is $\tau_{\gamma\gamma} \simeq 0.3 F_{X,-9} n_0^{1/2} t_3^{1/2} E_{52}^{-1/2} D_{28}^2 (\delta t / t) \epsilon_{X,\text{keV}}^{-1}$. Due to the low absorption depth and the flat spectral slope above $\epsilon_{IC,M}$, TeV photons associated with bright X-ray flares are expected to be detectable with detectors such as MAGIC, Milagro, H.E.S.S., VERITAS and ARGO etc.

THE ANISOTROPIC IC SCATTERING EFFECT

The incoming X-ray flare photons, which moves out in radial direction, are anisotropic seen by the isotropically distributed electrons in the forward shock, so more head-on scatterings may decrease the IC emission in the $1/\Gamma$ cone along the direction of the photon beam (corresponding to the angle less than $\pi/2$ relative to the photon beam direction in the comoving frame) [11], but enhance the emission at larger angles, with about a fraction of $3/8$ of total emission falling into the angles between $1/\Gamma$ and $2/\Gamma$ [7]. In the comoving frame of the forward shock, the IC emission power depends on the relative angle θ_s between the scattered photon direction and the seed photon beam direction as $P(\theta_s) \propto (1 - \cos\theta_s)^2$ [8]. The factor $(1 - \cos\theta_s)^2$ results in more power emitted at large angles relative to the seed photon beam direction. By simple algebraical calculation, one can obtain the fraction of photons scattered into the angles $0 \leq \theta_s \leq \pi/2$ in the shock comoving frame (corresponding to the $1/\Gamma$ cone of the relativistic afterglow jet in the observer frame due to aberration of light), i.e. $f_s \simeq \int_0^{\pi/2} (1 - \cos\theta_s)^2 \sin\theta_s d\theta_s / \int_0^\pi (1 - \cos\theta_s)^2 \sin\theta_s d\theta_s = 1/8$. Similarly, one can get that the fraction of photons falling into the cone of $2/\Gamma$ (corresponding to $\theta_s = 2\text{arctg}2$) is $1/2$. So the effect of anisotropic scattering can decrease the IC emission in the $1/\Gamma$ cone along the direction of the photon beam, but enhance the emission at larger angles. However if the opening angle is $\theta_j \gg 1/\Gamma$, the jet geometry can be approximately regarded as a sphere, and the observed IC power after integration over angles should be the same in every direction in the observer frame [9, 6]. Supposing that the flare photons and forward shock ejecta have a 4π solid angle (i.e. not jet), then the IC emission in the observer frame should have the same flux in every direction with a flux level as estimated in the above section. Hence the received IC component fluence is not reduced.

The anisotropic scattering effect does affect the temporal behavior of the IC emission. IC emission will be lengthened by the forward shock angular spreading time, which is also influenced by the anisotropic IC effect, so the IC component would not be correlated strictly with the flare light curves. Since a large fraction of the IC emission is falling within angles θ_{IC} between $1/\Gamma$ to $2/\Gamma$, the IC component will peak at

$$t_{IC,p} \sim R/2\Gamma^2 c + R(1 - \cos\theta_{IC})/c \sim R/2\Gamma^2 c + R\theta_{IC}^2/2c \sim R/\Gamma^2 c \sim 2t_{\text{flare},p} \quad (3)$$

where $t_{\text{flare},p} \sim R/2\Gamma^2 c$ is the flare peak time. The IC component light curves is shown as the thick solid line in Fig.2. The self-IC of X-ray flares may also produce a high-energy component, but its light curve should be correlated with the x-ray flare itself to some extent [6, 12], as shown in the thin solid line. GLAST might be able to distinguish these two different processes in future with the temporal behavior of the very high-energy emission and therefore provides a useful approach for distinguishing whether X-ray flares originate from late central engine activity or from external shocks.

The enhanced cooling of the forward shock electrons by the X-ray flare photons may suppress the synchrotron emission of the afterglows during the flare period[6]. Due to more head-on scatterings with electrons moving in angles larger than $\pi/2$ relative to the photon beam direction in the comoving frame and less scatterings with electrons within $\pi/2$, the anisotropic scattering effect may weaken the cooling effect of flare photons on the afterglow electrons (which are moving within $\pi/2$ in the comoving frame and




FIGURE 2. A schematic cartoon for the light curves of the X-ray flares (lower panel) and the predicted IC emission (upper panel) at GeV-TeV energy from different processes. In the upper panel, the solid line represents the IC emission of X-ray flare photons scattered by afterglow electrons, while the thick solid line denotes the self-IC emission of flare photons in the context of internal-shock or external-shock scenarios for X-ray flares.

moving within $1/\Gamma$ in the observer frame correspondingly), hence reduce the extent to which the flare photons affect the afterglow flux intensity. To suppress the synchrotron afterglow emission significantly, the cooling induced by the flare photons should be higher than that by the magnetic field. As a result, only bright flares are expected to significantly suppress the optical afterglow flux during the flare period, such as seen in GRB050904 [13].

REFERENCES

1. Burrows, D. et al. 2005a, *Science*, 309, 1833
2. Beloborodov, A. M. 2005, *ApJ*, 618, L13
3. Falcone, A. D. et al., 2006, *ApJ*, 641, 1010
4. Zhang, B. et al. 2006, *ApJ*, 642, 354
5. O'Brien P. T. et al. 2006, *ApJ*, 647, 1213
6. Wang, X. Y., Li, Z. & Mészáros, P. 2006, *ApJ*, 641, L89
7. Wang, X. Y. & Mészáros, P. 2006, *ApJ*, 643, L95
8. Rybicki, G. B.; Lightman, A. P., 1979, *Radiative Processes in Astrophysics* (New York: Wiley), p200
9. Fan, Y. Z. & Piran, T. 2006, *MNRAS*, 370, L24
10. Hurley, K. et al., 1994, *Nature*, 371, 652
11. Ghisellini, G. et al. 1991, *MNRAS*, 248, 14
12. Galli, A. et al. 2007, this proceedings
13. Gou, L. J. , Fox, D. & Mészáros, P. 2006, *ApJ*, astro-ph/0612256.